

AD-A162 590

DIFFERENTIATION OF CATALYTIC ACTIVITY OF SACCHAROMYCES  
CEREVISIAE CYTOCHR. (U) DEFENCE RESEARCH ESTABLISHMENT  
SUFFIELD RALSTON (ALBERTA) A R BHATTI ET AL. OCT 85  
DRES-SM-1148

1/1

UNCLASSIFIED

F/G 6/1


NL

DATE

END

10/11/85

DTA



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A



National Defence  
Défense nationale

UNCLASSIFIED



**DRES**

**SUFFIELD MEMORANDUM**

NO. 1148

**DIFFERENTIATION OF CATALATIC ACTIVITY OF *SACCHAROMYCES CEREVISIAE* CYTOCHROME B<sub>2</sub>: L-LACTATE DEHYDROGENASE FROM CATALASE T BY STARCH GEL ELECTROPHORESIS (U)**

by

A.R. Bhatti, T.C.M. Seah\* and J.G. Kaplan\*\*

Project No. 16 A 10

DTIC  
ELECTE  
DEC 11 1985  
S D E

\* Department of Biology, University of Ottawa, Ottawa, Ontario  
K1N 6N5, Canada

\*\* Department of Biochemistry, University of Alberta, Edmonton,  
Alberta, T6G 2P5, Canada

October 1985

DEFENCE RESEARCH ESTABLISHMENT SUFFIELD, RALSTON, ALBERTA



Canada

**WARNING**

The use of this information is permitted subject to  
recognition of proprietary and patent rights.

This document has been approved  
for public release and sale; its  
distribution is unlimited.

85 12 10 026

DTIC FILE COPY

AD-A162 590

UNCLASSIFIED

DEFENCE RESEARCH ESTABLISHMENT SUFFIELD  
RALSTON, ALBERTA

SUFFIELD MEMORANDUM NO. 1148

DIFFERENTIATION OF CATALATIC ACTIVITY OF SACCHAROMYCES CEREVISIAE CYTOCHROME  
B<sub>2</sub>: L-LACTATE DEHYDROGENASE FROM CATALASE T BY STARCH GEL ELECTROPHORESIS (U)

by

A.R. Bhatti, T.C.M. Seah\* and J.G. Kaplan\*\*

ADDRESS: \*\* Department of Biochemistry, University of Alberta, Edmonton,  
Alberta, T6G 2P5, Canada

\* Department of Biology, University of Ottawa, Ottawa, Ontario,  
K1N 6N5, Canada

UNCLASSIFIED

This document has been approved  
for public release and sale; its  
distribution is unlimited.

(A)

UNCLASSIFIED

DEFENCE RESEARCH ESTABLISHMENT SUFFIELD  
RALSTON, ALBERTA

SUFFIELD MEMORANDUM NO. 1148

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

DIFFERENTIATION OF CATALATIC ACTIVITY OF SACCHAROMYCES CEREVISIAE CYTOCHROME  
B<sub>2</sub>: L-LACTATE DEHYDROGENASE FROM CATALASE T BY STARCH GEL ELECTROPHORESIS (U)

by



A.R. Bhatti, T.C.M. Seah and J.G. Kaplan

ABSTRACT

Crude extracts of yeast exhibit two catalase activity bands on starch gel zymograms. Antibody prepared against catalase T, specifically precipitated fast moving catalatic band of catalase T, but did not affect the slow moving catalatic band due to cytochrome b<sub>2</sub>. *Keywords: Catalase; yeast; proteins*

UNCLASSIFIED

UNCLASSIFIED

DEFENCE RESEARCH ESTABLISHMENT SUFFIELD  
RALSTON, ALBERTA

SUFFIELD MEMORANDUM NO. 1148

DIFFERENTIATION OF CATALATIC ACTIVITY OF SACCHAROMYCES CEREVISIAE CYTOCHROME  
B<sub>2</sub>: L-LACTATE DEHYDROGENASE FROM CATALASE T BY STARCH GEL ELECTROPHORESIS (U)

by

A.R. Bhatti, T.C.M. Seah and J.G. Kaplan

### INTRODUCTION

Some years ago, in the course of purifying a highly active and typical catalase (catalase T) in S. cerevisiae (22), we became aware of the presence, in the same crude extracts, of three other proteins which also possessed catalatic activity; two of these were then purified to homogeneity (23). One of these was an atypical catalase which we called catalase-A (24), while the second one, containing non-heme iron and RNA, we

UNCLASSIFIED

called the catalatic nucleoprotein (25). Purification and properties of these two proteins have been described (24, 25). Ruis and his associates have elucidated the biosynthesis of catalase T and catalase A in yeast by means of double labelling with L-[<sup>3</sup>H]-leucine and <sup>59</sup>Fe coupled with immunoprecipitation techniques (29). More recently, they have studied the regulation of these hemoproteins in yeast (8) and have isolated and cloned the gene for catalase T (27). This communication presents evidence to show that the fourth protein in the yeast crude extract is the L-lactate: ferricytochrome C oxidoreductase (EC 1.1.2.3) which is commonly referred to as the yeast cytochrome b<sub>2</sub> (1).

Catalase (H<sub>2</sub>O<sub>2</sub>:H<sub>2</sub>O<sub>2</sub> oxidoreductase EC 1.11.1.6) catalyzes the decomposition of hydrogen peroxide directly into molecular oxygen and water. This mode of reaction distinguishes it from peroxidase (donor: H<sub>2</sub>O<sub>2</sub> oxidoreductase EC 1.11.1.7) which can also utilize H<sub>2</sub>O<sub>2</sub> as substrate (12). The substrate specificity of catalase was made use of by Scandalios (18) to localize, and to identify the presence of catalase(s) on starch gel zymograms; this method has been widely used. While non-enzymatic destruction of H<sub>2</sub>O<sub>2</sub> by ferric salts and by a number of haemoproteins has long been known (15), the possibility that the very weak catalatic activity of these metallic salts and proteins might give a positive reaction that might be misinterpreted for the presence of catalase on the zymogram has not been considered. There is now a voluminous literature (2, 5-7, 9, 13, 14, 18-21) dealing with catalase isoenzyme demonstrated by means of starch gel zymograms of crude extracts of a variety of species. We show in this communication that one of the band in a "catalase" zymogram of yeast crude extracts is due to the L-lactate:ferricytochrome C oxidoreductase (EC 1.1.2.3) which is commonly referred to as yeast cytochrome b<sub>2</sub>(1). Workers in this field should be aware of this, in order to avoid possible misinterpretation of their data.

## MATERIALS AND METHODS

Yeast cytochrome  $b_2$ , L-lactate dehydrogenase (LDH), was purchased from Worthington (Freehold, New Jersey) and yeast catalases were purified from commercial yeast (Standard Brands) as described (1, 3, 4). Antibody to purified catalase T was prepared by subcutaneously injecting, in the thigh, two 2-3 month old male rabbits with 1.0 mL of an enzyme preparation containing 4.0 mL of pure antigen (approximately 3.0 mg protein) and an equal volume of Freund's (complete) adjuvant. Three injections were given to each rabbit at intervals of one week and blood was collected from the ear at 3 to 8 days after the last injection. The  $\gamma$ -globulin fraction was precipitated from pooled sera by the method of Keckwick (11), resuspended in phosphate buffer ( $5 \times 10^{-3}$  M, pH 7.0) and dialyzed for 48 h at 4°C against the same buffer. The resulting preparation was used as the anti-catalase T serum.

The titanium sulfate method for catalase activity is expressed as the pseudo-first order rate constant per second per mg protein (22). Yeast cytochrome  $b_2$  activity was assayed by the method of Appleby and Morton (1) and specific activity is expressed as units per mg protein where one unit of activity is equal to the reduction of one micromole of ferricyanide per minute at 25°C. In some experiments, the catalatic activity of cytochrome  $b_2$  was assayed in the presence of  $1 \times 10^{-5}$  to  $1 \times 10^{-1}$  M 3-amino-1,2,4-triazole (AT). A modified method of Heim, Appleman, and Pyfrom (4) for catalase inhibition in vitro was employed. The method was as follows; one mL portions of enzyme solution containing a final concentration of  $5 \times 10^{-3}$  M phosphate buffer (pH 7.0),  $1 \times 10^{-3}$  M ethylenediaminetetraacetic acid (EDTA), 5 mg enzyme and a given concentration of AT, in the absence of  $H_2O_2$ , were incubated at 37°C for  $\frac{1}{2}$  hour after which the residual catalase and LDH activities were determined. Starch gel electrophoresis was carried out as previously described (3). Electrophoresis was performed at 4°C for



4 to 5 hours at 200 volts. The gel was sliced and the catalase activity was localized by a modification of the method of Scandalios (18). The sliced gels were incubated at room temperature in  $1 \times 10^{-1}$  M  $H_2O_2$  in  $5 \times 10^{-3}$  M phosphate buffer (pH 7.0). After 10 min, gels were rinsed thoroughly with distilled water and treated for 10 to 15 sec with a solution of 15% potassium iodide (KI) prepared in 5% acetic acid. The KI solution was poured off as soon as the background of the gels turned pale blue and the gels were rinsed thoroughly in a continuous stream of tap-water. The gels appeared dark blue except for the clear band where catalase activity was localized. The stained gels were preserved in a solution of ethanol, glycerol, and water (v/v 5:1:5). Protein concentration was determined by Lowry's method (17) with bovine serum albumin as standard.

### RESULTS AND DISCUSSIONS

The cytochrome  $b_2$  used in the present study was found to be homogeneous by analytical ultracentrifugation. By our method of assay, its catalatic activity was approximately 0.1% of that of the purified yeast catalase T (Table 1). Nonetheless, its catalatic activity was higher by an order of magnitude than that of the other haemoproteins which we have examined (Table 1). On starch gel zymograms, we found that 50  $\mu$ g of this purified enzyme was sufficient to give an unmistakable catalase activity band. Fresh crude extracts of yeast always exhibit two catalase activity bands on starch gel zymograms, one fast moving and one slow moving band towards the anode (Figure 1, sample 3). Of the two catalase activity bands, the fast moving one was due to yeast catalase T, since it was removed from the zymogram when the crude extract was first mixed with an excess of antibody to catalase T (Figure 1, sample 4). This was confirmed by using pure antigen, in the presence and absence of antibody to catalase T (Figure 1, sample 1 and 2), where sample 1 contained purified catalase T

and sample 2 contained catalase T after treatment with antibody to catalase T. The slow moving activity band was found to have an identical Rf value to that of purified yeast cytochrome  $b_2$  on the starch gel zymogram (Figure 1, sample 6). Pretreatment of cytochrome  $b_2$  with antibody to catalase T did not remove its catalatic activity (Figure 1, sample 7), showing that the catalatic activity of the yeast cytochrome  $b_2$  was not due to catalase T.

With aged crude extracts or extracts prepared from cells pretreated with n-butanol, three activity bands were observed. The third activity band moved between the fast and slow bands, slightly ahead of the slow moving band. This catalase activity band may be due to a proteolytic degradation (10) product of cytochrome  $b_2$ . We obtained identical results with crude extracts prepared from a variety of laboratory wild-type haploid and diploid strains of S. cerevisiae. Antibody to catalase T reacted specifically to its antigen and did not cross-react with cytochrome  $b_2$ . This was true in studies utilizing immunoprecipitation coupled with enzyme assay, micro-immunodiffusion and the standard Ouchterlony procedure.

The zymogram did not detect the presence of catalase A or catalatic nucleoprotein despite their relatively high specific activities, presumably due to their low concentrations in the crude extracts.

In separate studies, pretreatment of purified yeast LDH with antibody to catalase A did not remove the slow moving catalase activity band on starch gel zymogram (data not shown). Amino triazole (AT) is a specific inhibitor of catalase (4). The catalatic activity of cytochrome  $b_2$  was completely inhibited by  $1 \times 10^{-1}$  M AT while it had a little effect on the LDH activity (Figure 2). At this concentration of AT, yeast catalase was inhibited by 70%. With crystalline beef liver catalase, 50% inhibition was obtained with  $5 \times 10^{-4}$  M AT.

Weak as was its catalatic activity compared to that of catalase T, yeast cytochrome  $b_2$  could be mistaken as a catalase isoenzyme on the starch gel zymogram. It is possible that similar non-catalase proteins which could catalyze the decomposition of  $H_2O_2$  may also be present in crude extracts of organisms other than yeast; thus starch gel catalase zymograms of cell extracts should be interpreted with caution. In the absence of more conclusive data, catalase activity bands demonstrated by starch gel zymogram are, at best, descriptive, showing the presence of proteins which can catalyze the decomposition of  $H_2O_2$  as opposed to a conclusive evidence for the presence of catalase(s) (28). For the same reason, it is equally misleading to assume that catalactic activity detectable by means of enzyme assay in a given crude extract preparation is due to the presence of catalase, particularly when the catalatic activity is low.

#### CONCLUSION

Precipitous readings of simple biochemical tests can frequently lead to erroneous conclusions. In this study, crude extracts of yeast were shown to exhibit two bands of apparent catalase activity on starch gel zymograms. One band was found to be due to catalase whereas the other was due to cytochrome  $b_2$  activity. It is possible that similar non-catalase proteins, capable of catalyzing the decomposition of  $H_2O_2$ , could also be present in the crude extracts of organisms other than yeast. Therefore, starch gel catalase zymograms of such cell extracts should be always interpreted with caution.

REFERENCES

1. Appleby, C.A. and R.K. Morton. 1959. Lactic dehydrogenase and cytochrome  $b_2$  of bakers yeast: Purification and crystalization. *Biochem. J.* 71: 492-499.
2. Beckman, L., J.G. Scandalios and L.B. Baker. 1964. Catalase hybrid enzymes in maize. *Science* 146: 1174-1175.
3. Bhatti, A.R. and J. Done. 1974. On the presence of two types of alkaline phosphatase in Serratia marcescens. *Microbios* 11: 203-213.
4. Heim, W.G., D. Applemann and H.T. Pyfrom. 1956. Effect of 3-amino-1, 2,4-triazole (AT) on catalase and other compounds. *Amer. J. Physiol.* 186: 19-23.
5. Holmes, R.S. 1972. Catalase multiplicity in normal and acatalaemic mice. *FEBS Lett.* 24: 161-164.
6. Holmes, R.S. and C.J. Masters. 1970. Epigenetic interconversions of multiple forms of mouse liver catalase. *FEBS Lett.* 11: 45-48.
7. Holmes, R.S. and C.J. Masters. 1972. Species specific features of the distribution and multiplicity of mammalian liver catalase. *Arch. Biochem. Biophys.* 148: 217-223.
8. Hortner, H., G. Ammerer, E. Hartter, B. Hamilton, J. Rytka, T. Bilinski and H. Ruis. 1982. Regulation of synthesis of catalases EC-1.11.1.6 and iso 1 cytochrome C in Saccharomyces cerevisiae by glucose oxygen and heme. *Eur. J. Biochem.* 128: 179-184.

9. Jones, G.L. and C.J. Masters. 1972. On the differential inhibition of the multiple forms of catalase EC-1.11.1.6 in mouse tissue. FEBS Lett. 21: 207-210.
10. Kaplan, N.O. 1968. Nature of multiple forms of enzymes. Ann. N.Y. Acad. Sci. 151: 382-399.
11. Keckwick, R.A. 1940. The serum proteins in multiple myelomatosis. Biochem. J. 34: 1248-1257.
12. Loew, O. 1901. Catalase, a new enzyme of general occurrence with special reference to the tobacco plant. U.S. Dept. Agr. Report 68: 47p.
13. Matsubara, S., H. Suter and H. Aebi. 1966. Heterogeneity of catalase in blood of heterozygous cases of acatalasia. Experientia. 22: 428-429.
14. Matsubara, S., H. Suter and H. Aebi. 1967. Fractionation of erythrocyte catalase from normal, hypo-catalatic and A-catalatic. Humangenetik. 4: 29-41.
15. Nicholls, P. and G.R. Schonbaum. 1963. "Catalases" in The Enzymes. 2nd ed., Ed. P. Boyer, H. Lardy and K. Mayback, New York: Academic Press Vol. 8: 150-225.
16. Numachi, K.I. 1971. Electrophoretic variants of catalase in the black rockfish. Bull. Jap. Soc. Sci. Fish, 37: 1177-1181.
17. Lowry, O.H., N.J. Rosenbrough, A.L. Farr and R.J. Randall. 1951. Protein measurement with Folin phenol reagent. J. Biol. Chem.

193: 265-275.

18. Scandalios, J.G. 1964. Tissue-specific isozyme variation in maize. J. Heredity. 55: 281-285.
19. Scandalios, J.G. 1965. Subunit dissociation and recombination of catalase isozymes. Proc. Nat. Acad. Sci. 53: 1035-1040.
20. Scandalios, J.G. 1968. Genetic control of multiple molecular forms of catalase in maize. Ann. N.Y. Acad. Sci. 151: 274-293.
21. Scandalios, J.G. 1969. Genetic control of multiple molecular forms of enzymes isozymes in plants. Biochem. Genet. 3: 37-79.
22. Seah, T.C.M. and J.G. Kaplan. 1973. Purification and properties of the catalase of Baker's yeast. J. Biol. Chem. 248: 2889-2893.
23. Seah, T.C.M., A.R. Bhatti and J.G. Kaplan. 1973. Abnormal minor catalatic proteins of yeast. Feb. Proc. 32: 1395.
24. Seah, T.C.M., A.R. Bhatti and J.G. Kaplan. 1973. Novel catalatic proteins of Baker's yeast. I. An atypical catalase. Can. J. Biochem. 51: 1551-1555.
25. Seah, T.C.M., A.R. Bhatti and J.G. Kaplan. 1978. A catalatic nucleoprotein of Baker's yeast. FEBS Lett. 85: 305-309.
26. Shabalina, A.T. and O.L. Serov. 1970. Genetics of blood catalase isoenzymes in Gallus domesticus. Dokl. Akad. Sel-skokhoz. Nauk. Bolg. 3: 369-372.

27. Spevak, W., F. Fessler, J. Rytko, A. Traczyk, M. Skoneczny and H. Ruis. 1983. Isolation of the catalase T structure gene of Saccharomyces cerevisiae by functional complementation. Molec. Cell. Biol. 3: 1545-1551.
28. Stavri, H. and D. Stavri. 1975. Electrophoretic mobility of the catalases of some mycobacteria. Arch. Roum. Path. Exp. Microbiol. 34: 85-59.
29. Zimniak, P., E. Hartter, W. Woloszczuk and H. Ruis. 1976. Catalase biosynthesis in yeast: Formation of catalase A and catalase T during oxygen adaptation of Saccharomyces cerevisiae. Eur. J. Biochem. 71: 393-398.

Figure 1

Starch gel zymogram showing catalase activity of purified yeast catalase T, crude extract of bakers' yeast and purified yeast cytochrome b<sub>2</sub>. Slots 1, 3, 6 contained purified catalase T (4 µg), a crude extract of bakers' yeast (125 µg) and purified yeast cytochrome b<sub>2</sub> (700 µg), respectively. Slots 2, 4, 7 were as in 1, 3, 6, except that the preparations had been treated with excess antibody to catalase T, incubated at 37°C for ½ hr, centrifuged and the clear supernatant used for electrophoresis. Slot 5 contained tracking dye.

Figure 2

The effect of 3-amino-1,2,4-triazole on the catalase and LDH activities of purified yeast cytochrome b<sub>2</sub>. Details in "Methods".






Figure 1

Starch gel zymogram showing catalase activity of purified yeast catalase T, crude extract of bakers' yeast and purified yeast cytochrome  $b_2$ . Slots 1, 3, 6 contained purified catalase T ( $4 \mu\text{g}$ ), a crude extract of bakers' yeast ( $125 \mu\text{g}$ ) and purified yeast cytochrome  $b_2$  ( $700 \mu\text{g}$ ), respectively. Slots 2, 4, 7 were as in 1, 3, 6, except that the preparations had been treated with excess antibody to catalase T, incubated at  $37^\circ\text{C}$  for  $1/2$  hr, centrifuged and the clear supernatant used for electrophoresis. Slot 5 contained tracking dye.




Figure 2

The effect of 3-amino-1,2,4-triazole on the catalase and LDH activities of purified yeast cytochrome  $b_2$ . Details in "Methods".

Table 1

Catalatic activity of non-catalase proteins

Enzymes	Specific activity K/sec/mg
Yeast cytochrome b <sub>2</sub>	$4.90 \pm 1.39 \times 10^{-4}$
Haemoglobin (beef blood)	$4.00 \pm 1.20 \times 10^{-5}$
Cytochrome c	no detectable activity
Horse radish peroxidase	no detectable activity
Haemin (non-protein)	no detectable activity
Yeast catalase T	$7.06 \times 10^{-1}$

## DOCUMENT CONTROL DATA - R &amp; D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall document is classified)

1. ORIGINATING ACTIVITY  DRES		2a. DOCUMENT SECURITY CLASSIFICATION Unclassified	
		2b. GROUP	
3. DOCUMENT TITLE DIFFERENTIATION OF CATALATIC ACTIVITY OF <u>SACCHAROMYCES CEREVISIAE</u> CYTOCHROME B <sub>2</sub> : L-LACTATE DEHYDROGENASE FROM CATALASE T BY STARCH GEL ELECTROPHORESIS			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
5. AUTHOR(S) (Last name, first name, middle initial)  A.R. Bhatti, T.C.M. Seah and J.G. Kaplan			
6. DOCUMENT DATE October 1985		7a. TOTAL NO. OF PAGES 12	7b. NO. OF REFS 29
8a. PROJECT OR GRANT NO.  16A10		9a. ORIGINATOR'S DOCUMENT NUMBER(S)  SM 1148	
8b. CONTRACT NO.		9b. OTHER DOCUMENT NO.(S) (Any other numbers that may be assigned this document)	
10. DISTRIBUTION STATEMENT  Unlimited			
11. SUPPLEMENTARY NOTES		12. SPONSORING ACTIVITY	
13. ABSTRACT  Crude extracts of yeast exhibit two catalase activity bands on starch gel zymograms. Antibody prepared against catalase T, specifically precipitated fast moving catalatic band of catalase T, but did not affect the slow moving catalatic band due to cytochrome b <sub>2</sub> .			

## KEY WORDS

Catalase, cytochrome B<sub>2</sub>, aminotriazole, Saccharomyces cerevisiae, electrophoresis, isoenzymes

## INSTRUCTIONS

1. **ORIGINATING ACTIVITY** Enter the name and address of the organization issuing the document.
- 2a. **DOCUMENT SECURITY CLASSIFICATION** Enter the overall security classification of the document including special warning terms whenever applicable.
- 2b. **GROUP** Enter security reclassification group number. The three groups are defined in Appendix 'M' of the DRB Security Regulations.
3. **DOCUMENT TITLE** Enter the complete document title in all capital letters. Titles in all cases should be unclassified. If a sufficiently descriptive title cannot be selected without classification, show title classification with the usual one-capital-letter abbreviation in parentheses immediately following the title.
4. **DESCRIPTIVE NOTES** Enter the category of document, e.g. technical report, technical note or technical letter. If appropriate, enter the type of document, e.g. interim, progress, summary, annual or final. Give the inclusive dates when a specific reporting period is covered.
5. **AUTHOR(S)** Enter the name(s) of author(s) as shown on or in the document. Enter last name, first name, middle initial. If military, show rank. The name of the principal author is an absolute minimum requirement.
6. **DOCUMENT DATE** Enter the date (month, year) of Establishment approval for publication of the document.
- 7a. **TOTAL NUMBER OF PAGES** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.
- 7b. **NUMBER OF REFERENCES** Enter the total number of references cited in the document.
- 8a. **PROJECT OR GRANT NUMBER** If appropriate, enter the applicable research and development project or grant number under which the document was written.
- 8b. **CONTRACT NUMBER** If appropriate, enter the applicable number under which the document was written.
- 9a. **ORIGINATOR'S DOCUMENT NUMBER(S)** Enter the official document number by which the document will be identified and controlled by the originating activity. This number must be unique to this document.
- 9b. **OTHER DOCUMENT NUMBER(S)** If the document has been assigned any other document numbers (either by the originator or by the sponsor), also enter this number(s).
10. **DISTRIBUTION STATEMENT** Enter any limitations on further dissemination of the document, other than those imposed by security classification, using standard statements such as:
  - (1) "Qualified requesters may obtain copies of this document from their defence documentation center."
  - (2) "Announcement and dissemination of this document is not authorized without prior approval from originating activity."
11. **SUPPLEMENTARY NOTES** Use for additional explanatory notes.
12. **SPONSORING ACTIVITY** Enter the name of the departmental project office or laboratory sponsoring the research and development. Include address.
13. **ABSTRACT** Enter an abstract giving a brief and factual summary of the document, even though it may also appear elsewhere in the body of the document itself. It is highly desirable that the abstract of classified documents be unclassified. Each paragraph of the abstract shall end with an indication of the security classification of the information in the paragraph (unless the document itself is unclassified) represented as (TS), (S), (C), (R), or (U).  
  
The length of the abstract should be limited to 20 single-spaced standard typewritten lines 7 1/4 inches long.
14. **KEY WORDS** Key words are technically meaningful terms or short phrases that characterize a document and could be helpful in cataloging the document. Key words should be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context.

**END**

**FILMED**

---

**1-86**

**DTIC**